

# Neurological Manifestations of HIV-AIDS among HIV diagnosed Patients at a Tertiary Care Hospital of Khyber Pakhtunkhwa, Pakistan

Yaseen Khan<sup>1</sup>, Syed Ather<sup>1</sup>, Amjad Taqveem<sup>1</sup>, Bilal Awan<sup>1</sup>, Aftab Khan<sup>2</sup>

Department of Medicine, Lady Reading Hospital<sup>1</sup>, Pakistan Health Research Council (PHRC) Research Centre<sup>2</sup>, Khyber Medical College, Peshawar.

## Abstract

**Background:** Neurological manifestations affecting the nervous system at all stages of Human Immunodeficiency Virus (HIV) infection are common. Neurological complications occur in more than 40% of patients with HIV infection. They are the presenting feature of Acquired Immune Deficiency Syndrome (AIDS) in 10-20% of cases. At autopsy the prevalence of neuropathological abnormalities is 80%. Although an ongoing decline in HIV associated Central Nervous System (CNS) diseases has been observed in very recent years, the mortality from these diseases remains high.

**Objective:** To study the type and frequency of different neurological involvements in patients with HIV infection at tertiary care hospital in Khyber Pakhtunkhwa Peshawar and to correlate them with CD4 counts.

**Study design, settings and Duration:** This retrospective observational study was carried at Lady Reading Hospital, Khyber Pakhtunkhwa, Peshawar, Pakistan over a period of 8 years from May 2009 to June 2017.

**Subjects and Methods:** A total of 100 HIV sero-positive patients of both genders, aged >18 years, showing clinical evidence of central nervous system (CNS) involvement were included. Their clinical manifestations, laboratory investigations, and neuroimaging were studied. Laboratory investigations along with magnetic resonance imaging (MRI), EEG and nerve conduction study of the brain/spine was also performed.

**Results:** Tuberculous meningitis was the most common presentation as secondary CNS illness (49%), followed by cryptococcal meningitis (16%) and cerebrovascular accidents (7%). Furthermore, 6% had neurosyphilis, 5% had acquired immune deficiency syndrome (AIDS) associated dementia and peripheral neuropathy occurred in 17% of the patients. Headache was the most common neurological symptom seen in 42% of the patients. Seizures were noted in 35% of the patients. CD4 was significantly low in most of the patients with progressive multifocal leukoencephalopathy, HIV associated encephalopathy (HAD) and cryptococcal meningitis compared with other neurological manifestations. CD4 counts in tuberculous meningitis and HIV associated encephalopathy were 115/ $\mu$ l and 83/ $\mu$ l, respectively.

**Conclusion:** CNS tuberculosis was the most common secondary infection seen in HIV patients followed by cryptococcal meningitis. A high index of clinical suspicion of neurological involvement in HIV patients helps in the early diagnosis and early institution of specific treatment, which in turn decreases the morbidity and mortality considerably. Early treatment and prophylaxis of neurological problems in HIV patients is very important to decrease the mortality rate.

**Key words:** Tuberculous meningitis, cryptococcal meningitis, HIV, neurological manifestation, CD4.

## Corresponding Author:

Yaseen Khan  
Department of Medicine, Lady Reading Hospital  
Peshawar.  
Email: [yaseen07k@gmail.com](mailto:yaseen07k@gmail.com)

**Received:** 16 November 2017, **Accepted:** 28 September 2018,  
**Published:** 29 September 2018

## Authors Contribution

YK conceptualized the project. SA & BA did the data collection. YK & AT performed the literature search. Statistical analysis was done by AK. YK, AT & AK also did the drafting, revision and writing of manuscript.

## Introduction

The causative agent of acquired immunodeficiency syndrome (AIDS) is human immunodeficiency virus (HIV). An estimated 37 million persons worldwide are infected with HIV. The main cause of its spread in both men and women is heterosexual mode.<sup>1</sup>

In Pakistan the HIV epidemic is still in infant stage and according to UNAIDS fact sheet 2016 report there are about 130,000 to 150,000 people are affected. In Khyber Pakhtunkhwa (KP), the high

rate of HIV infection is because of male sex workers, unsafe blood transfusion, and irrational use of injection practices.<sup>2</sup> Additionally, other factors like illiteracy, overseas employment in gulf, poverty, lack of sex education, high prevalence of intravenous drug users and high prevalence of sexually transmitted diseases (STDs) particularly in transgender are the contributing factors in the transmission of HIV.<sup>3</sup>

In Pakistan, the HIV patients have higher chances of progressing to full-blown AIDS and developing neuro-tuberculosis, because patients either do not seek medical advice because of social stigma, do not take antiretroviral treatment (ART) or become ART defaulters. The AIDS epidemic in KP Pakistan is still in its infancy.<sup>4</sup> However, HIV-positive individuals may become significant threat in the coming years, and will affect the planned national economy and health setup.

The nervous system is extensively involved in HIV and no part of nervous system being immune to the virus. Therefore, the knowledge about central nervous system (CNS) complications of HIV is very important for medical professionals. However, limited studies are available regarding the neurological manifestations of HIV/AIDS in Pakistan. Despite the high prevalence rate of HIV neurological manifestations, not a single study on this topic has been conducted in Khyber Pakhtunkhwa, Peshawar Pakistan.

## Subjects and Methods

The study was retrospective cross sectional study conducted over a period of 8 years from May 2009 to June 2017 at a tertiary care Hospital, Government Lady Reading Hospital Peshawar Pakistan. Total 100 HIV patients with neurological manifestations were included. HIV Patients with neurological manifestation having incomplete data were excluded from the study. It is the main tertiary care hospital of KP Province, which also caters for FATA and adjoining parts of Afghanistan. A detailed history was noted in each patient with special emphasis on history of sexual contacts, drug abuse, travel abroad, penile or vaginal ulcers, surgical interventions, blood transfusions or a history of tuberculosis, recurrent herpes zoster, seizures, unconsciousness or headache. Detailed physical examination was carried out in each patient to look for opportunistic infections. Fundus was examined in each patient. Routine biochemical, hematological, and bacteriological testing including liver and kidney function tests, complete blood count, cerebrospinal fluid (CSF) examination (cytology, sugar, proteins), special staining studies (Ziehl–Nelson stain, Indian ink, etc.), chest X-ray, and (cryptococcal and

toxoplasmosis antigen with specific antigen detection tests in the serum and VDRL) were performed. CD4 count was done for all the study participants. All patients underwent reactive fluorescent treponemal antibody tests. Magnetic resonance imaging (MRI), EEG and nerve conduction study of the brain/spine were also performed.

Every patient's information was kept highly confidential. Patients with a diagnosis of neuro-AIDS were treated according to WHO guidelines<sup>5,6</sup> using first line antiretroviral treatment (ART), antituberculous, antifungal medications, antiviral, antibiotics, and anticonvulsants and anti-psychiatric medications. Associated opportunistic infections were treated with respective drugs. Intensive care unit management and mechanical ventilation were provided whenever required. Patients having CD4 count <500cells/ $\mu$ L were put on Highly Active Antiretroviral Therapy (HAART) as recommended.<sup>5,6</sup> Chemoprophylaxis and ART categories was instituted as indicated. Trimethoprim–sulfamethoxazole was given to prevent pneumonia in all patients with a CD4 lymphocyte count of <200cells/ml. Fluconazole and azithromycin was used for fungal and MAC complex prophylaxis. Data was entered and analyzed using SPSS version 16.

## Results

A total of 100 patients were admitted with neurological manifestations of HIV/AIDS. The mean age of patient was  $36\pm 19$  (range 18-55 years). Male to female ratio was 2:1. Thirty one (31%) patients had a primary neurological illness, whereas 69 (69%) patients had secondary illnesses (Table-1).

**Table 1: Age and gender wise distribution of study population.**

| Particulars | Patients (n= 100) |
|-------------|-------------------|
| Age         |                   |
| 18 – 30     | 22                |
| 31 – 40     | 45                |
| 41 – 50     | 25                |
| Above 50    | 08                |
| Gender      |                   |
| Male        | 75                |
| Female      | 25                |

Out of 100 patients with a neurological manifestation, 59 (59%) presented with fever. Headache was reported in 42 patients (42%), while 35 patients (35%) presented with a history of convulsions. Altered sensorium was the presenting complaint in 62 patients (62%). Vomiting was present in 35 (35%) patients. Twenty-seven (27%) patients presented with tingling and numbness. Forty-one

(41%) patients presented with focal deficits. Bladder and bowel involvement was found in 40 (40%) patients. Signs of meningeal irritation were present in 55 (55%) patients (Table-2).

**Table 2: Clinical manifestations in the presence of neurological illness in HIV positive patients.**

| Clinical presentation         | No. of cases | %  |
|-------------------------------|--------------|----|
| Fever                         | 59           | 59 |
| Altered consciousness         | 62           | 62 |
| Headache                      | 42           | 42 |
| Convulsion                    | 35           | 35 |
| Tingling and numbness         | 27           | 27 |
| Vomiting                      | 35           | 35 |
| Forgetfulness                 | 12           | 12 |
| Bowel and bladder impairment  | 40           | 40 |
| Focal neurological deficit    | 41           | 41 |
| Signs of meningeal irritation | 55           | 55 |

Result showed that majority of the females (78%) acquired infection from their HIV positive husbands and 65% among total number of patients acquired infection sexually. The study provided that 08% patients acquired HIV infection through blood or blood related products and 18.5% patients were found infected through intravenous drug use (IDU). About 4.5% of the patients were infected through homosexual route, while in 4.15% patients the cause of HIV infection was unidentified.

**Table 3: Primary and secondary neurological illness observed in HIV positive patients.**

| Neurological Illness | Types   | (n=100)<br>(%) |
|----------------------|---|----------------|
| Primary              | Distal symmetric polyneuropathy (DSPN)          | 17             |
|                      | AIDS dementia complex (ADC)                     | 05             |
|                      | Acute inflammatory demyelinating polyneuropathy | 02             |
|                      | Stroke syndromes                                | 02             |
| Secondary            | Tuberculous bacterial meningitis                | 49             |
|                      | Cryptococcal meningitis                         | 16             |
|                      | Toxoplasmosis                                   | 03             |
|                      | Progressive multifocal leukoencephalopathy.     | 03             |
|                      | neurosyphilis                                   | 06             |

Distal symmetrical polyneuropathy (DSPN) was found in 17% HIV patients. Cerebrovascular disease was detected in 7% of HIV infected individuals. Present study provided that tuberculous meningitis (TBM) as found in 49% of HIV patients and considered common secondary CNS illness in the present study, followed by cryptococcal meningitis (16%). Toxoplasmosis was observed in 3%, progressive multifocal leukoencephalopathy in 3% and neuro-syphilis in 6% of the cases. Three patient had herpes simplex encephalitis (presenting

with altered sensorium and seizures with bilateral orbital and temporal lobe involvement) and 3 patients had cerebral toxoplasmosis which presented with right-sided partial seizure with secondary generalization. One patient had herpes zoster, affecting T4-t7 thoracic dermatome on the right side, presenting with radicular lancinating pain (Table-3).

HIV patients with CNS infections presented with low levels of CD4 counts i.e. <200/μL. The average CD4 count in cells/mm<sup>3</sup> in the presence of various clinical presentations was as follows: 170±100.9 in DSPN, 245.7±113.9 in AIDP, 120.3±81.8 in tuberculous meningitis, 41.0±29.60 in cryptococcal meningitis, and 55.6±38.7 in toxoplasmosis (Table-4). In HIV-associated dementia (HAD), the mean CD4 count was 85±81.72. Statistically, there was no difference in the ART status and the development of neurological manifestations in HIV/AIDS patients.

**Table 4: Mean CD4 count in neurological illness in HIV positive patients.**

| Neurological illness                            | Mean CD4 Count ± Standard Deviation (SD) |
|---|--|
| Tuberculous bacterial meningitis                | 120.3±81.8                               |
| Cryptococcal meningitis                         | 41.0±29.60                               |
| Toxoplasmosis                                   | 55.6±38.7                                |
| Progressive multifocal Leucoencephalopathy      | 115±75.5                                 |
| Neurosyphilis                                   | 125±45.5                                 |
| HIV-associated dementia (HAD)                   | 85±81.72                                 |
| Distal symmetric polyneuropathy                 | 170±100.9                                |
| Acute inflammatory demyelinating Polyneuropathy | 245.7±113.9                              |

## Discussion

HIV-infected patients having CNS complication can be divided into meningitis, encephalopathy intracerebral space occupying lesions, and spinal cord process. Where effective ART treatment implemented, these complications have declined markedly. As the age of HIV, the cognitive declines have been observed. In the hippocampus, basal ganglia, prefrontal cortex, and white matter slow neuro-degradation occur due to infected macrophages infiltrate the brain parenchyma. Release of IL-1, TNF, and IL-6 causes low damage to the CNS.<sup>8</sup>

In the present study, the age group 24-47 was found with high grade of neurological involvement, which correlates with a study done by Sircar et al.<sup>9</sup> The age group 24-27 is a highly productive section of our society put alarming effect in growth of a developing country.

Heterosexual transmission of HIV disease was also observed in this study. The study provided

similar pattern of results like studies from India and other Asian countries. While in Europe homosexual transmission is a commonest route.

In this study ratio of Male to female was 2:1, which is comparable to the study done by Soluet al and Sircar et al from India.<sup>9,10</sup> Tuberculous involvement was seen in 49 (49%) patients, out of which 09 had intracranial tuberculomas and 40 had tuberculous meningitis. This is similar to the findings of other studies in which the incidence of tuberculous involvement was 47% and 42%, respectively.<sup>11,12</sup> In tuberculous meningitis common presenting symptoms were fever (55%) followed by headache (35%), altered consciousness (32%), and convulsions (12%).

It was observed that in United States, 5-10% patients had AIDS with cryptococcal meningitis the most common problem in HIV patients.<sup>13</sup> In present study, 16 out of 100 (16%) patients with neuro-AIDS had cryptococcal meningitis. The other symptoms like headache, fever and vomiting were the complaints in HIV patients with neurological manifestations. In HIV patients' Cryptococcal neoformans causes minimal inflammation with defused immune mechanism, which explains why neck rigidity and photophobia were infrequent in this study. Cryptococcal meningitis occurs in terminally ill patients with CD4 count <100  $\mu\text{L}$ .<sup>14</sup> CSF examination of cryptococcal meningitis patients showed the average protein levels of 132%, sugar of 62%, and total cell of 45/mm<sup>3</sup> and lymphocytes of 95%. CSF cryptococcal antigen titer was used to diagnose in this study. Other studies documented the similar pattern of results as reported in previously.<sup>15</sup> The incidence of mortality due to cryptococcal meningitis in patients with HIV infection was reported in 45%. The poor prognostic predictors were altered mentation at the time of diagnosis, positive Indian ink staining, positive blood culture, a low CSF leukocyte count, high cryptococcal antigen titers, and increased CSF opening pressure. PML, Cryptococcal meningitis, toxoplasmosis and neurosyphilis reported were similar to the findings of two other studies by Abayomi et al and Rana et al.<sup>16,17</sup> In a Brazilian study and another by Deshpande et al., toxoplasmosis was considered main cause of CNS complication.<sup>18,19</sup> The most sensitive technique MRI was used to diagnose this entity. This infection generally occurred in HIV patients with low CD4 count.<sup>20</sup>

Progressive multifocal leuko encephalopathy (PML) was developing in 4% of AIDS patients, and it was main complication of AIDS in 25% of HIV cases. However, in the present study, 15 patients had such kind of history.

Neuro-syphilis a main complication and its association is not unexpected because both AIDS

and neuro-syphilis are sexually transmitted diseases. But neuro-syphilis develop in late stages of disease.<sup>21</sup> Invasion of CNS by the organism *Treponema palladium* can also be expected to occur. Patients with neuro-syphilis had serum VDRL titers  $\geq 1:16$ . The median of serum CD4 cell and VDRL I counts were 127cell/ $\mu\text{L}$  and 1:122 respectively.

HIV-associated myelopathy, neurocognitive impairment, myopathy, aseptic meningitis and peripheral neuropathy are primary illnesses. Another study revealed same kind of information as documented in this study.<sup>22</sup> Eight patients (8%) were diagnosed to be having a cerebrovascular accident (CVA). Five of them had an infarct in the basal ganglia and other three had an infarct in the cerebellum. CVA evidence provided in the present study is similar to that seen in another study.<sup>23</sup> DSPN is mostly seen in patients with low CD4 counts. DSPN is also the most common side-effects of ART. ADC was seen in 7% of our patients, which was diagnosed by the mini mental score examination (MMSE). Neuro-cognitive dysfunction generally occurred when less than 100 cell/ $\mu\text{L}$  CD4 count was reported.<sup>24</sup>

Mean CD4 count (in cell/ $\mu\text{L}$ ) in this study was 151 $\pm$ 100.9 with DSPN, 259.7 $\pm$ 113.9 with AIDP, 120.3 $\pm$ 81.8 with TBM, 42.0 $\pm$ 29.60 with cryptococcal meningitis, and 54.6 $\pm$ 38.7 with toxoplasmosis. A study conducted by Deshpande from India was similar with this study.<sup>25</sup> In HAD, the mean CD4 count was 91 cells/ $\mu\text{L}$ , which is comparable with the NIMS study in which mean CD4 count was 92 cells/ $\mu\text{L}$ .<sup>26</sup> HIV patients with CD4 cell count below 200 are at a high risk of developing serious illnesses, which has been seen in this study too. The presence of neurological manifestations is associated with decreased CD4 counts. Cryptococcal meningitis was associated with lowest CD4 counts, and had a fatal outcome in 40% of the patients, followed by toxoplasmosis, HAD and neurosyphilis (Table-5). In KP Pakistan today, the only affordable methodology is the estimation of the CD4 cell counts. It can serve as a guide to assess the HIV status, assess the risk for the development of neurological manifestations in such patients and help in instituting timely intervention in the form of prophylaxis and treatment.<sup>27</sup> The mortality was 25% in TBM, 15% in toxoplasmosis and 40% in cryptococcal meningitis (Table-4). In the era of HAART and specific treatment that is available for opportunistic infections, mortality in patients with HIV with neurological illness has considerably decreased. Lady Reading hospital, being KP biggest territory hospital most patients of HIV/AIDS are referred for treatment and management. The study provided useful information that neurological problems should be identified and

treated at early stage of HIV to decrease the morbidity and mortality.

The reported CNS opportunistic infections in HIV patients in this study provided the need of intervention on neurological manifestation in HIV patients. A proper guideline of treatment and diagnosis of CNS manifestation should be a part of ART treatment. Awareness is needed among health professionals regarding neurological manifestation among HIV patients to address the issue before entering HIV patient in a terminal stage of diseases.

### Acknowledgement

The Authors are grateful to the staffs of Lady reading hospital, HIV center Hayatabad medical complex and PHRC Center Khyber Medical Collage for their consent to carry out this research and their timely support.

**Conflict of interest:** None declared.

### References

1. HIV/AIDS. Fact sheet No. 360, Geneva, Switzerland: World Health Organization; (Accessed on 15th June 2016) Available from URL:<http://www.who.int/media/centre/factsheets/fs360/en/>
2. Correa M, Gisselquist D. HIV from blood exposures in India-an exploratory study. Colombo: Norwegian Church Aid, South Asia, Colombo, 2005.
3. Malani PN. Human immunodeficiency Virus. JAMA Patient Page 2016; 316(2): 238.
4. National Survey on HIV. Published in daily Dawn 1ST October 2017. Accessed on 15th June 2018) Available from URL:<https://www.dawn.com/news/1361153>
5. Solu MD, Dandge V, Karla P. Clinical profile of HIV patients with neurological manifestation. Guj Med J 2012; 67(2): 98-103.
6. CDC Global HIV/AIDS Milestones: On the path to an AIDS-Free Generation. 2011. Accessed on 15th June 2018) Available from URL:<https://www.cdc.gov/globalaids/Global-HIV-AIDS-at-CDC/AIDS-Free-generation.html>
7. Patil VC, Patil HV. Neurological manifestations of HIV-AIDS at a tertiary care centre in western Maharashtra. Int J Med Public Health 2014; 4: 210-17.
8. Shripad RK, Annad C, Pramod J. Neurological manifestations in HIV positive patients attending tertiary care teaching hospital in Western India. Int J Adv Med 2016 3(4): 1055-8.
9. HIV fact sheets 2003-2006. National AIDS Control Organization. Ministry of Health and Family Welfare, November 2007, Government of KP Pakistan.
10. Sircar AR, Tripathi AK, Choudhary SK, Misra R. Clinical profile of AIDS: A study at a referral hospital. J Assoc Physician India 1998; 46: 775-8.
11. Solu MD, Dandge V, Kalra P, Prajapati R, Josh A. Clinical profile of HIV positive patients neurological manifestations. Guj Med J 2012; 67(2): 98-103.
12. National strategies and algorithm on HIV testing (algorithm III). Guidelines on HIV testing. NACO March 2007, Ministry of Health and Family Welfare.
13. Das CP, Sawhney IMS. Neurological complications of HIV infection. Neurol Indian J 1998; 46: 82-93.
14. Fauci AS, Lane HC. Human immunodeficiency virus disease: Harrison's Principle of Internal Medicine. Vol 1. 17th ed. New York: Mcgraw Hill; 2008. pp. 1139-83.
15. Wadhwa A, Kaur R, Bhalla P. Profile of central nervous system disease in HIV/AIDS patients with special reference to cryptococcal infection. Neurologist 2008; 14: 247-51.
16. Rana HM, Doshi DA, Virpariya KM, Shah AN, Somani SS. A study of clinical profile of HIV positive patients with neurological manifestations. HIV AIDS Rev 2011; 10: 76-9.
17. Millogo A, Ki-Zerbo GA, Sawadogo AB, Ouedraogo I, Yameogo A, Tamini MM, et al. Neurologic manifestations associated with HIV infections at the Bobo-Dioulasso Hospital Center (Burkina Faso). Soc Pathol Exot 1999; 92: 23-6.
18. Rana HM, Doshi DA, Virpariya KM, Shah AN, Somani SS. A study of clinical profile of HIV positive patients with neurological manifestations. HIV & AIDS Review 2011; 171-4.
19. Oliveira JF, Greco DB, Oliveira GC, Christo PP, Guimaraes MD, Oliveira RC. Neurological disease in HIV infected patients in the era of highly active anti-retroviral treatment: A Brazilian experience. Rev Soc Bras Med Trop 2006; 39: 146-51.
20. Teja VD, Talasila SR, Vemu L. Neurologic manifestations of HIV infection: An Indian hospital based study. AIDS Read 2005; 15: 139-43.
21. Abayomi OS, Ojini F, Okubadejo N, Danesi NM, Kalapo K, Osalusi B, et al. Pattern and outcome of neurological manifestations of HIV/AIDS- A review of 154 cases in a Nigerian university teaching hospital- A preliminary report. African J Neurol Sci 2005; 24: 29-36.
22. Kovacs JA, Kovacs AA, Polis M, Wright WC, Gill VJ, Tuazon CU. Cryptococcosis in acquired immunodeficiency syndrome. Ann Intern Med 1985; 103: 533-8.
23. Attili VS, Gulati AK, Singh VP, Varma DV, Rai M, Sundar S. Neurological manifestations in HIV infected patients around Varanasi, KP Pakistan. BMC Infect Dis 2006; 6: 39.
24. Sing R, Kaur M, Arora D. Neurological complications in late-stage hospitalized patients with HIV disease. Ann Indian Acad Neurol 2011; 14: 172-7.
25. Deshpande AK, Patnaik MM. Non-opportunistic neurologic manifestations of the Human immunodeficiency virus: An Indian Study. J Int AIDS Soc 2005; 7: 2.
26. Brightbill TC, Ihmeidean IH, Donovan Post MJ, Berger JR, Katz DA. Neurosyphilis in HIV-positive and HIV negative patients: Neuroimaging findings. AJNR Am J Neuroradiol 1995; 16: 703-11.
27. Mc Arthur JC. Neurologic manifestations of AIDS. Medicine 1987; 66: 407-37.